

NSF-ITP-92-107

YCTP-P20-92

Spontaneous CP Violation Small Instanton & Invisible Axion

Soo-Jong Rey*

Institute for Theoretical Physics

University of California, Santa Barbara CA 93106

&

Center for Theoretical Physics, Sloane Laboratory

Yale University, New Haven CT 06511 USA

September 1992

Abstract

I propose a variant invisible axion model of spontaneous CP violation at the electroweak scale without CP domain-wall and ‘strong CP’ problems. Both large sized QCD and small sized non-QCD instantons break CP and Peccei-Quinn symmetries, and render cosmologically harmful CP domain walls unstable. The decaying epoch depends on size of small instanton effects, and is around $100eV$ if the current neutron electric dipole moment bound is maximally saturated. The model satisfies constraints from primordial D and He photo-dissociation and black hole formation, while producing cosmologically interesting size of gravitational waves and galaxy-scale density perturbations.

*Yale-Brookhaven SSC Fellow. Bitnet: Soo@Yalehep. This work was supported in part by Texas National Research Laboratory Commission.

1. Motivation:

Origin of CP violations[1] still remains elusive. Current experimental data from $K^o - \bar{K}^o$ system and neutron electric dipole moment (NEDM):

$$\epsilon \leq 2.26 \cdot 10^{-3}; \quad \left| \frac{\epsilon'}{\epsilon} \right| \leq 1.45 \cdot 10^{-3}; \quad \bar{\theta}_{\text{qcd}} \leq 10^{-13} \left(\frac{m_N}{m_q} \right). \quad (1)$$

indicate *strong violation* of weak, flavor nondiagonal CP (FNCP) symmetry and *weak violation* of strong, flavor diagonal CP (FDCP) symmetry.

Complex Yukawa coupling constants or complex Higgs coupling constants are known to provide sources of explicit CP violations. Alternatively, CP violations may arise spontaneously[2] [3] during the electroweak symmetry breaking, for example, as in extended technicolor theories[4]. In addition, in supersymmetric standard models, spontaneous CP violation offers an attractive solution[5] of suppressing other potentially large contributions to NEDM than $\bar{\theta}_{\text{qcd}}$. It has been known, however, that this option is cosmologically disastrous due to appearance of stable CP domain walls in the early universe[6]. The walls quickly dominate energy density of the universe, and lead too short the present age to our Universe.

Recently, I proposed a particle physics solution to this cosmological CP domain wall problem[7], utilizing nonperturbative QCD effect. The idea was as follows. QCD instanton effect is practically negligible around the electroweak scale at which CP is spontaneously broken. However, the effect becomes pronounced as the QCD scale $\Lambda_{\text{qcd}} \sim 0.2\text{GeV}$ is approached and induces operators

$$\mathcal{L}_{\text{nonp}} = K_{\text{qcd}} e^{i\bar{\theta}_{\text{qcd}}} (\bar{U}_L U_R \bar{D}_L D_R \bar{S}_L S_R) + h.c. \quad (2)$$

Under CP transformation,

$$\begin{aligned} \bar{\theta}_{\text{qcd}} &\rightarrow -\bar{\theta}_{\text{qcd}} \\ (CP)\mathcal{L}_{\text{nonp}}(CP)^{-1} &\neq \mathcal{L}_{\text{nonp}}. \end{aligned} \quad (3)$$

Unless $\bar{\theta}_{\text{qcd}}$ is 0 or π , CP, a discrete global symmetry, is explicitly broken by the QCD instantons. Being a nonperturbative effect, size of the intrinsic CP violation is tiny; $K_{\text{qcd}} \sim \Lambda_{\text{qcd}} m_u m_d m_s \ll M_W^4$. This lifts the vacuum degeneracy connecting two sides of the CP domain wall, and leads to a subsequent wall annihilation driven by a false vacuum decay.

A drawback of this idea was the ‘strong-CP’ problem in assuming $\bar{\theta}_{\text{qcd}}$ nonzero but small $\sim 10^{-9}$ to meet Eq.(1). One may introduce Kim’s[8] (KSVZ) or Dine-Fischler-Srednicki[9] (DFSZ) invisible axion as a viable solution to the ‘strong-CP’ problem. However, in the same paper[7], I also have shown that spontaneous CP violation and the Peccei-Quinn (PQ) symmetry are not compatible each other, at least, in the simplest two-Higgs model. It was found that there always exists a nonanomalous discrete symmetry, which is a simultaneous CP and finite PQ transformations:

$$D : (PQ)_{2\bar{\theta}}(CP) \mathcal{L} (CP)^{-1} (PQ)_{2\bar{\theta}}^{-1} = \mathcal{L}. \quad (4)$$

Thus, there remains yet another cosmological domain wall problem when the new discrete symmetry D is spontaneously broken at the electroweak scale. Furthermore, especially in DFSZ axion model, I also observed that the PQ symmetry drives the Higgs potential minima to a CP-conserving point at least in the simplest two electroweak Higgs model.

In this paper, I propose a viable solution to these difficulties, and make PQ symmetry compatible with spontaneous CP violation at the electroweak scale. I achieve this by (1) extending the electroweak Higgs sector and (2) taking into account of additional nonperturbative effects from non-QCD sector, for example, technicolor gauge group or semiclassical gravity. Point (1) allows a nontrivial EW vacuum in which CP is spontaneously broken, while point (2) lifts a vacuum degeneracy connected by the discrete symmetry of Eq.(4). Furthermore, the unstable but long-lived domain wall may serve an extra source of long-wavelength gravitational waves and galaxy scale density perturbations.

2. Model:

I first state criteria what a viable model is required to meet:

- (1) *intrinsically broken strong CP and an invisible axion solution to the ‘strong CP’ problem;*
- (2) *spontaneously broken weak CP during the electroweak phase transition.*

(Our viewpoint is different from Nelson-Barr[10] in that flavor-nondiagonal CP begins to appear at the electroweak scale, and that ‘strong-CP’ problem is solved by an invisible axion.)

I introduce a singlet Higgs field Σ and add one more to weak doublet Higgs

fields H_i in the standard axion models. Thus, the standard KSVZ model is now extended to Lee's two-Higgs doublet model[2], while the standard DFSZ model to Weinberg's three-Higgs doublet model[3]. We do not, however, introduce Glashow-Weinberg (GW) criterion[11] of natural flavor conservation (NFC); approximate flavor symmetry for small Yukawa coupling constants is already technically natural[12]. The GW criterion usually imposes an additional discrete symmetry to the Yukawa and Higgs sector, which is spontaneously broken at the electroweak symmetry breaking. Cosmologically, this causes yet another domain wall problem, which I wanted to avoid.

Σ field acquires a vacuum expectation value (VEV) $\langle \Sigma \rangle \approx f_a \sim 10^{12} \text{ GeV}$, and breaks PQ symmetry spontaneously. Radiative corrections do respect PQ and CP symmetries as exact symmetries. Thus, effective electroweak Higgs potential near the weak scale consists of all renormalizable gauge, PQ and CP invariant operators among Higgs doublets H_i and axion $a \sim \arg \Sigma$. Schematically, it reads

$$V_{\text{Higgs}}[H, \Sigma] = \sum_{ij} M_{ij}(\Sigma) H_i^\dagger H_j + \sum_{ijkl} \lambda_{ijkl}(\Sigma) (H_i^\dagger H_j)(H_k^\dagger H_l) + \sum_{ijkl} \tilde{\lambda}_{ijkl}(\Sigma) (H_i^\dagger \vec{\tau} H_j)(H_k^\dagger \vec{\tau} H_l). \quad (5)$$

Hermiticity and functional dependence of M^2 , λ , $\tilde{\lambda}$ to the axion field Σ is fixed by CP and PQ invariances of the low-energy Higgs potential V_{Higgs} . Low-energy Yukawa couplings are similarly constrained. Minimization of the above electroweak Higgs potential was analyzed previously[3][7]. For both Lee-KSVZ and Weinberg-DFSZ models, it can be shown that there does exist a region of parameter space in which the CP and gauge symmetries are spontaneously broken: $\langle \phi_a \rangle = \frac{v_a}{\sqrt{2}} e^{i\theta_a}$ with $\theta_a - \theta_b \neq n\pi$. (Related multi-Higgs models were studied previously[13].)

In addition, QCD and non-QCD instantons generate additional terms to the electroweak Higgs potential. An example of non-QCD nonperturbative effect is the gravitational instanton. The simplest gravitational instanton with an asymptotically flat spacetime is the Eguchi-Hanson metric[14]. The EH instanton does not break the PQ symmetry. On the other hand, a charged EH instanton can do. An EH instanton of size ρ with a self-dual electromagnetic

field (EM instanton) is known[14]:

$$g^{\mu\nu} = \delta^{\mu\nu} - \frac{\rho^4}{x^4} \frac{x^\mu x^\nu}{x^2} + \frac{\rho^4}{x^4 - \rho^4} \frac{\tilde{x}^\mu \tilde{x}^\nu}{x^2}; \quad A_\mu = 2P \frac{\rho^2 \tilde{x}^\mu}{x^4} \quad (6)$$

in which $x^2 \equiv x^\mu x^\mu \geq \rho^2$ and $\tilde{x}^\mu = (y, -x, t, -z)$ is identified with x^μ because of $\mathbf{RP}_3$ spacetime topology. The instanton EM charge $P \in 3\mathbf{Z}$ to give a compatible spin structure for the standard model chiral fermions. Instanton action is found $S_{\text{EM}} = \frac{\pi P^2}{\alpha_e(\rho)} \gg 1$, hence, semiclassical approximation is valid. In the QCD and EM instantons backround, the PQ current is anomalous:

$$\nabla^\mu J_{\mu\text{pq}} = \sum_{\text{ferm}} Q_{\text{pq}} \left(\frac{2Q_{\text{em}}^2}{16\pi^2} F_{\mu\nu} \tilde{F}_{\mu\nu} + \frac{2C_2(\mathbf{R})}{8\pi^2} \text{Tr} \mathbf{G}_{\mu\nu} \tilde{\mathbf{G}}_{\mu\nu} \right) \quad (7)$$

in which the sum is over all fermions with nontrivial PQ charge. In the dilute instanton approximation, the instantons induce multi-fermion operators similar to Eq.(2). They constitute the *only* sources of intrinsic PQ and CP symmetry breaking (potentially dangerous irrelevant operators). Tying up fermion zero modes with PQ and CP conserving Yukawa and Higgs self-interaction vertices, a low-energy Higgs-axion potential is found;

$$V_{\text{inst}} = \int_{f_a^{-1}}^{\Lambda_{\text{qcd}}^{-1}} d\mu[\rho] [K_{\text{qcd}}(\rho) e^{i\bar{\theta}_{\text{qcd}} \Sigma^m} + K_{\text{em}}(\rho) e^{i\bar{\theta}_{\text{em}} \Sigma^n} + h.c.]. \quad (8)$$

Magnitude of EM instanton induced axion potential K_{em} was estimated in detail[15] to give $K_{\text{em}} \sim 10^{-9} K_{\text{qcd}}$ in which $K_{\text{qcd}} \sim 10^{-6} \text{GeV}^4$. Thus, modifications in the above electroweak Higgs potential minimization due to QCD and EM instantons are totally negligible. The CP-odd coupling constants $\bar{\theta}$'s do receive finite renormalizations across each fermion mass thresholds and from the spontaneous CP violating phases θ_{ab} . Instanton multiplicity $m, n \in \mathbf{Z}_+$ does depend on details of axion model. Since each term in Eq.(8) comes from different instanton effect, I expect $m \neq n$ in general. QCD and EM instantons give:

$$m \equiv \text{Tr} 2C_2(R); \quad n \equiv \text{Tr} 18Q_{\text{em}}^2 \quad (9)$$

in which trace is over all fermions with nonzero PQ charge assignment.

Each of the operators appearing in Eq.(8) breaks CP and PQ symmetries separately. If either K_{qcd} or K_{em} were absent, the combined Higgs potential $V_{\text{Higgs}} + V_{\text{inst}}$ is manifestly invariant under simultaneous transformations of CP:

$\bar{\theta} \rightarrow -\bar{\theta}$ and a finite PQ: $arg\Sigma \rightarrow arg\Sigma - 2/m$ (or $2/n$) $\cdot \bar{\theta}$. This is the new discrete symmetry D alluded earlier in Eq.(4), and is the origin of another cosmological domain wall problem.

Important point is that this ceases to be the case if *both* K 's are present. CP transformation changes:

$$\bar{\theta}_{\text{qcd}}, \bar{\theta}_{\text{em}} \rightarrow -\bar{\theta}_{\text{qcd}}, -\bar{\theta}_{\text{em}}. \quad (10)$$

Thus, *no* subsequent finite PQ transformation

$$arg\Sigma \rightarrow \Sigma - \alpha \quad (11)$$

can rotate both $\bar{\theta}$'s back to the original values *simultaneously*. Since PQ symmetry is the only other available symmetry to rotate phases in V_{inst} , I draw a conclusion that there simply does not exist any residual, nonanomalous discrete symmetry such as D in Eq.(4).

The size of additional instanton induced operator, K_{em} is further constrained. Minimizing the axion potential Eq.(8), I find

$$m \langle a \rangle + \bar{\theta}_{\text{qcd}} \approx \frac{K_{\text{em}}}{K_{\text{qcd}}} \cdot \left(\frac{n^2}{m^2} \bar{\theta}_{\text{qcd}} - \frac{n}{m} \bar{\theta}_{\text{em}} \right). \quad (12)$$

Taking $|\bar{\theta}_{\text{qcd}} - \bar{\theta}_{\text{em}}| \sim \mathcal{O}(\pi)$, this implies

$$K_{\text{em}} \leq 10^{-9} K_{\text{qcd}} \quad (13)$$

to be consistent with NEDM measurement Eq.(1). As obvious as it may sound, however, nonzero value of $\bar{\theta} \equiv m \langle a \rangle + \bar{\theta}_{\text{qcd}} (1 \text{ GeV}) \sim 10^{-9}$ makes the CP domain wall unstable as well!

Energy density difference between the otherwise degenerate vacua connected by D symmetry is estimated from the potential Eq.(8):

$$\Delta\mathcal{E}_{\text{vac}} \approx K_{\text{em}} \sin 2\left(\frac{n}{m} \bar{\theta}_{\text{qcd}} - \bar{\theta}_{\text{em}}\right) \sin(n \langle a \rangle + \bar{\theta}_{\text{em}}). \quad (14)$$

In case Eqs. (12) and (13) are maximally saturated,

$$\Delta\mathcal{E}_{\text{vac}} \approx 10^{-9} K_{\text{em}} \approx (100 \text{ eV})^4. \quad (15)$$

It is utterly small compared to the QCD scale, yet large enough to be cosmologically significant as shall be discussed below.

3. Cosmological Evolution of Unstable Domain Wall:

Cosmological evolution of the CP domain wall was studied in my previous work[7]. At the epoch PQ symmetry is spontaneously broken, the QCD and EM instanton induced potential Eq.(6) is negligible. This remains true even until an electroweak transition temperature T_{EW} is reached and the CP domain walls are formed. As the temperature goes down further, energy density of the practically stable domain walls starts to grow faster than that of the background radiation. Due to background plasma viscosity, the domain wall is straightened out only to a scale $R_c(T) \sim \sqrt{M_w^3 M_{pl}}/T^3$, which is a fraction $F \equiv R_c(T)H(T)$ of the horizon at a given temperature T . Using this, epoch that energy density of the wall begins dominate over that of radiation is found as[7]:

$$\rho_{\text{wall}} \approx \rho_{\text{rad}} \quad \rightarrow \quad T_c \approx 10^{-8} \frac{1}{\sqrt{\lambda}} T_{EW} \quad (16)$$

most conservatively, in which λ denotes a typical value of Higgs quartic couplings λ_{ijkl} and $\tilde{\lambda}_{ijkl}$. With $\lambda \sim 10^{-4}$, the domain wall starts to dominate energy density of the universe only below $T_c \approx 300 \text{eV}$. In particular, this implies that primordial nucleosynthesis (PNS) at the MeV scale is not disturbed by the presence of CP domain walls, as their energy density is completely negligible at the PNS epoch.

The otherwise stable CP domain wall begins to feel the energy density difference in Eq.(8) as the pressure provided by the vacuum energy density difference begins dominate over the viscosity $\Delta\mathcal{E}_{vac} \geq T^4$. This occurs when

$$T^* \approx 10^{-6} \Lambda_{\text{qcd}} \approx 100 \text{ eV}. \quad (17)$$

The wall then moves quickly to the speed of light via false vacuum decay. As the wall reaches the speed of light, regions of false vacuum are quickly driven away.

That $T^* \leq \frac{1}{3}T_c$ implies a brief period of domain-wall dominated universe. This period is prolonged if the EM instanton size K_{em} is smaller than the estimate I used; $\sim 10^{-9}K_{\text{qcd}}$. This transient, domain-wall dominated epoch near T^* is after the present day galaxy sized regions first came inside the horizon around $T \sim 1 \text{KeV}$. Relativistic motion of the walls tend to wash out primordial density fluctuations smaller than the horizon scale, i.e., the galaxy scale.

Instead, randomly located walls generate a Poisson distribution of density perturbations at scales smaller than the present galaxy size. Magnitude of this perturbation is therefore expected to be $\delta\rho/\rho \sim (R_c(T)H)^{1/2} \approx F^{1/2}$. Furthermore, as the domain wall dominated epoch gives rise to a modest power-law inflation $H(t) = 2/t$, it is expected that the density perturbation of galaxy size is further damped, and may provide a ‘biasing’ mechanism to the standard galaxy formation.

Violent wall collision will produce both strong metric perturbation which in turn forms gravitational waves (GW) and primordial black holes (PBH). Since the walls move quickly with a relativistic velocity, time scale of collision is expected to be $\sim R_c(T^*)$. Thus radiated gravitational wave energy E_{gw} and wavelength $\lambda(T^*)$ with a quadrupole Q is estimated:

$$E_{\text{gw}} = \frac{1}{M_{\text{pl}}^2} \left(\frac{d^3 Q}{dt^3} \right)^2 \approx \frac{1}{M_{\text{pl}}^2} \frac{E_v^2}{R_c}; \quad \lambda(T^*) \sim R_c(T^*) \quad (18)$$

in which $E_v = \frac{4\pi}{3} R_c^3 \Delta\mathcal{E}_{\text{vac}}$. Therefore, we find $\Delta E_{\text{gw}}/E_v \approx F^2$ and is about 15–25% at $T = T^*$. Thus, a significant fraction of the false vacuum energy goes into the gravitational wave radiation[16]. Using the fact that the gravitational wave vector $2\pi/\lambda$ and amplitude $h(\lambda)$ both scales as $\frac{1}{R(T)}$, I finally estimate them at present time as:

$$h(\lambda) \approx 10^{-8}; \quad \lambda \approx 10^{23}(\text{cm}). \quad (19)$$

Both the amplitude and frequency fall short an order of magnitude of interesting ranges that millisecond pulsar and large scale cosmic microwave background radiation experiments[17] may be able to detect! Uncertainties in the above rough estimate should be reflected in Eq.(19); a better quantitative estimate might be able to put the present model against upcoming cosmological experimental tests.

Similarly, PBH’s may form as the wall collide. With the wall-induced density fluctuation $\delta\rho/\rho$ estimated to $\sim F^{1/2} \sim 50 - 60\%$ I expect the PBH formation inefficient even at the locus of wall collisions. Typical mass of PBH’s is estimated heavy enough $\sim \epsilon T^{*4} \cdot (R_c^2/H) \approx \epsilon F^2(T^{*4}/H^3)$ that its Hawking radiation is completely negligible. Here, ϵ denotes a geometrical factor for the locus of wall collisions at which the PBH’s are formed, and is expected to be

$\epsilon \ll 1$. Maximal fraction of the PBH formation energy ΔE_{pbh} is constrained, in case the practically stable PBH's dominate the matter of the present universe, by $\Delta E_{\text{pbh}}/E_v \leq (13.6\text{eV}/T^*) \sim 13\%$. This bound is viewed large enough to accomodate even the most efficient PBH formation possible. Correspondingly, PBH produced gravitational waves are negligible compared to the one generated by wall collision alluded above.

The wall collision also produces $\mathcal{O}(\frac{1}{\lambda})$ energetic Higgs H^* and subsequently cascading lighter particles including photons γ^* , to which PNS puts a strong constraint. The energetic photons may photo-dissociate and change D and He abundances from the successfully predicted values of the standard PNS model. However, the energetic particles scatter predominantly with the 10^9 more abundant, cosmic background photons γ_{cb} than D's and He's. This will continue until $T \sim 10^3 \text{ eV}$ [18] at which number of the high-energy tail γ_{cb} 's is comparable to the number of D and He's, and threshold of $H^* + \gamma_{\text{cb}}$ or $\gamma^* + \gamma_{\text{cb}} \rightarrow l\bar{l}$ is reached. This is precisely the scale that CP domain walls completes decaying. Thus, PNS constraint of D and He abundances is seen marginally compatible with the present model.

I thank L.M. Krauss, L. McLerran and M. Srednicki for useful discussions. I also acknowledge hospitality of the Institute for Theoretical Physics at Santa Barbara where this research was finished. This work was supported in part by the National Science Foundation under grant No. PHY89-04035, and by DOE and TNRLC.

References

- [1] For an comprehensive collection of original papers, see L. Wolfenstein, *CP Violation*, North-Holland Pub. Co. (1989).
- [2] T.D. Lee, Phys. Rev. **D8** (1973) 1326; Phys. Rep. **9** (1974) 143, 176; see also, J. Liu and L. Wolfenstein, Nucl. Phys. **B289** (1987) 1.
- [3] S. Weinberg, Phys. Rev. Lett. **37** (1976) 657; see also, G. Branco, Phys. Rev. **D22** (1980) 2901.
- [4] E. Eichten, K. Lane and J.P. Preskill, Phys. Rev. Lett. **45** (1980) 225.

- [5] N. Maekawa, Phys. Lett. **282B** (1992) 387; A. Pomarol, *Spontaneous CP Violation in Supersymmetric Theories*, SCIPP-92/30 preprint (1992).
- [6] Ya.B. Zel'dovich, I.B. Kobzarev and L. Okun, Phys. Lett. **B50** (1974) 340.
- [7] L.M. Krauss and S.-J. Rey, Phys. Rev. Lett. (1992).
- [8] J.E. Kim, Phys. Rev. Lett. **43** (1979) 103; M.A. Shifman, V.I. Vainshtein and V.I. Zakharov, Nucl. Phys. **B166** (1980) 493.
- [9] M. Dine, W. Fischler and M. Srednicki, Phys. Lett. **109B** (1981) 199; A.P. Zhitniskii, Sov. J. Nucl. Phys. **31** (1980) 260.
- [10] A. Nelson, Phys. Lett. **136B** (1984) 387; S.M. Barr and A. Zee, Phys. Rev. Lett. **55** (1985) 2253; S.M. Barr, Phys. Rev. **D34** (1986) 165.
- [11] S.L. Glashow and S. Weinberg, Phys. Rev. **D15** (1977) 1958.
- [12] A. Antaramian, L.J. Hall and A. Rasin, *Flavor Changing Interactions Mediated by Scalars at the Weak Scale*, LBL-32345 preprint (1992).
- [13] A.R. Zhitniskii, Yad. Fiz. **31** (1980) 1024 [Sov. J. Nucl. Phys. **31** (1980) 529]; C.Q. Geng, X.D. Jiang and J.N. Ng, Phys. Rev. **D38** (1988) 1628.
- [14] T. Eguchi and A. Hanson, Phys. Lett. **74B** (1978) 430; Ann. Phys. (NY) **120** (1979) 82; G. 't Hooft, Nucl. Phys. **B315** (1989) 517; see also, S.-J. Rey, *Gravitational Superstring Instantons and Solitons*, NSF-ITP-92-106, YCTP-P18-92 preprint (1992).
- [15] R. Holman, T.W. Kephart and S.-J. Rey, *Semiclassical Gravity and Invisible Axions*, NSF-ITP-92-85, YCTP-P17-92 preprint (1992).
- [16] M.S. Turner and F. Wilczek, Phys. Rev. Lett. **65** (1990) 3080.
- [17] D.R. Stinebring *et al.*, Phys. Rev. Lett. **65** (1990) 285.
- [18] D. Lindley, Mon. Not. R. Astron. Soc. **188** (1979) 15P; **193** (1980) 593; L.M. Krauss, Phys. Rev. Lett. **53** (1984) 1976; S. Sarker and A.M. Cooper, Phys. Lett. **148B** (1984) 347.